

Preprint SSU-HEP-97/03
Samara State University

**CONTRIBUTION OF HADRONIC VACUUM
POLARIZATION TO HYPERFINE
SPLITTING OF MUONIC HYDROGEN**

Faustov R.N.

Scientific Council "Cybernetics" RAS,
117333, Vavilova, 40, Moscow, Russia,

Martynenko A.P.

Samara State University, 443011, Ac. Pavlov, 1, Samara, Russia

Abstract

The contribution of hadronic vacuum polarization to the hyperfine splitting of the muonic hydrogen ground state is calculated with the account of experimental data on the cross section of $e^+e^- \rightarrow$ annihilation into hadrons and the dipole parameterization of electromagnetic proton form factors.

Over many years the comparison of experimental and theoretical results for hyperfine splitting of hydrogen, muonium and positronium ground state energy level is considered as an important verification of our understanding of the bound state problem in quantum electrodynamics. Especially high precision was achieved at the present time when measuring hydrogen atom hyperfine structure [1, 2]:

$$\Delta E_{exp.}^{hf.s} = 1420405.7517667(9) KHz. \quad (1)$$

Theoretical value of hyperfine splitting, which ordinary written in the form

$$\Delta E_{th.}^{hf.s} = \Delta E^F (1 + \delta^{QED} + \delta^S + \delta^P), \quad \Delta E^F = \frac{8}{3} \alpha^4 \frac{\mu_P m_P^2 m_e^2}{(m_P + m_e)^3}, \quad (2)$$

(μ_P is the proton magnetic moment, expressed in nuclear magnetons; m_P , m_e are proton and electron masses correspondingly), contains besides electromagnetic correction δ^{QED} some more two terms δ^S and δ^P . These corrections take into account the influence of strong interaction. The term δ^S describes the effects of proton finite size and recoil contributions, and δ^P is the correction of proton polarizability. Particle structure members δ^S and δ^P are absent for purely leptonic systems [3]. Electrodynamical correction δ^{QED} for hyperfine structure of hydrogen atom, accounting relativistic and radiative effects, is known at present time with the accuracy 0,12 ppm [4]. The increase of accuracy in theoretical expression (2) is impossible without the detailed consideration of strong interaction in the terms δ^S and δ^P . The calculation of different contributions to the hydrogen atom hyperfine splitting (HFS), connected with proton structure (correction δ^s) was done in the paper [4]. Authors of [5] have proposed an experimental method to obtain more exact information about proton electromagnetic structure. It based on the measurement of hyperfine splitting of the muonic hydrogen (μp) ground state with the accuracy 10^{-4} . It is the authors opinion that such measurement will allow to find the relation between the corrections δ^S and δ^P , what will increase the accuracy of theoretical formula (2). In the case of muonic hydrogen, when the light electron must be replaced by the heavy muon, the relative contribution of different effects in (2) is changed. So, the hadronic vacuum polarization contribution, which was omitted in (2) in view of it's small value by comparison with δ^S and δ^P , may have more relative contribution for muonic hydrogen HFS. The effect of hadronic vacuum polarization must be considered in order to extract correctly the parameters δ^S and δ^P from experiment. In this paper we have calculated the hadronic vacuum polarization contribution to the hyperfine structure of muonic hydrogen on the

basis of quasipotential method [6]. In this approach the particle interaction operator for 1γ - and 2γ - processes takes the form [7]:

$$V = V_{1\gamma} + V_{2\gamma} = V^c + \Delta V, \quad (3)$$

$$V_{1\gamma} = T_{1\gamma}, \quad V_{2\gamma} = T_{2\gamma} - T_{1\gamma} \times G^f \times T_{1\gamma},$$

where V^c is Coulomb potential, $T_{1\gamma}$, $T_{2\gamma}$ are one- and two-photon scattering amplitudes beyond energy surface, $[G^f]^{-1} = (b^2 - \vec{p}^2)/2\mu_R$ is the free two-particle Green function (μ_R is relativistic reduced mass, b^2 is the square of particle relative motion momentum on energy surface). Then we can write the displacement of energy level as follows:

$$\Delta E = \langle \psi^c | \Delta V | \psi^c \rangle, \quad (4)$$

where ψ^c is the ordinary Coulomb wave function. The hadronic vacuum polarization contribution to the energy spectrum is determined by Feynman diagrams represented on Fig.1 by means (3). To take into account vacuum polarization in these one-loop diagrams, we must do the following substitution in photon propagator [8]

$$\frac{1}{k^2 + i\varepsilon} \rightarrow \left(\frac{\alpha}{\pi}\right)^2 \int_{s_{th.}}^{\infty} \frac{\rho(s) ds}{k^2 - s + i\varepsilon}, \quad (5)$$

where the spectral function $\rho(s)$ is connected with known cross section of e^+e^- - annihilation to hadrons σ^h :

$$\rho(s) = \frac{R(s)}{3s} = \frac{\sigma^h(e^+e^- \rightarrow hadrons)}{3s\sigma_{\mu\mu}(e^+e^- \rightarrow \mu^+\mu^-)}, \quad (6)$$

and $\sigma_{\mu\mu}(e^+e^- \rightarrow \mu^+\mu^-) = 4\pi\alpha^2/3s$ is the e^+e^- - annihilation cross section to muonic pair.

Let consider the proton factor, corresponding to two-photon exchange diagrams:

$$M_p^{\mu\nu} = \frac{\Gamma^\mu(\gamma^0 E_2 - \hat{p} + m_p)\Gamma^\nu}{p^2 + b^2 - 2E_2 p_0 + i\varepsilon} + \frac{\Gamma^\nu(\gamma^0 E_2 + \hat{p} + m_p)\Gamma^\mu}{(p+t)^2 + b^2 + 2E_2 p_0 + i\varepsilon}, \quad (7)$$

where p^μ is the loop 4-momentum, 4-momentum $t = (0, \vec{p} - \vec{q})$ is determined by relative motion momenta $\vec{p}$, $\vec{q}$ of initial and final states correspondingly. Proton vertex function Γ^μ may be expressed through Dirac $\rho(p^2)$ and Pauli $f(p^2)$ form factors, describing the proton electromagnetic structure:

$$\Gamma^\mu = \left[\gamma^\mu \rho(p^2) + \frac{i}{2m_p} \sigma^{\mu\nu} p_\nu f(p^2) \right], \quad (8)$$




Figure 1: Feynman diagrams, defining the hadronic vacuum polarization contribution to hyperfine structure of atom (μp)

where $\rho(0) = e$, $f(0) = e\kappa$, κ is the proton anomalous magnetic moment: $\kappa=1.792847386(63)$ [1]. Muonic tensor of considered one-loop amplitudes is equal:

$$M_{muon}^{\mu\nu} = \frac{\gamma^\mu(\gamma^0 E_1 + \hat{p} + m_p)\gamma^\nu}{p^2 + b^2 + 2E_1 p_0 + i\varepsilon}. \quad (9)$$

The hadronic vacuum polarization contribution to the ground state hyperfine splitting of muonium and positronium was obtained in [9]. It is clear that for exact calculation of discussed one-loop corrections in (μp), we must consider the dependence of form factors $\rho(p^2)$, $f(p^2)$ from loop momentum p . To do numerical calculations of the corrections $O(\alpha^6)$ caused by hadronic vacuum polarization we can use dipole parameterization of proton electromagnetic form factors of the form:

$$\rho(p^2) = \frac{G_E - \frac{p^2}{4m_p^2}G_M}{1 - \frac{p^2}{4m_p^2}}, \quad f(p^2) = \frac{G_M - G_E}{1 - \frac{p^2}{4m_p^2}}, \quad (10)$$

$$G_M = \frac{1 + \kappa}{\left(1 - \frac{p^2}{\Lambda^2}\right)^2}, \quad G_E = \frac{1}{\left(1 - \frac{p^2}{\Lambda^2}\right)^2}, \quad (11)$$

where the parameter of proton structure was taken in the kind $\Lambda = 0, 898m_p$ [1]. Multiplying the expressions (7) and (9) for construction of quasipotential, we obtain the following sum of form factor terms with the accuracy $O(\alpha^6)$:

$$\begin{aligned}
\langle M_p^{\mu\nu} M_{muon}^{\mu\nu} \rangle = & -\rho^2(p^2) \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \left(\frac{1}{D_p(-p)} + \frac{1}{D_p(p)} \right) \frac{1}{D_\mu(p)} - \\
& -2p^2 \frac{\rho(p^2) f(p^2)}{m_p} \langle \gamma^\mu \gamma^\nu \rangle \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \left(\frac{1}{D_p(-p)} + \frac{1}{D_p(p)} \right) \frac{1}{D_\mu(p)} + \\
& +4\rho(p^2) f(p^2) \langle \hat{p} \gamma^\mu \rangle \langle \hat{p} \gamma^\mu \rangle \left(\frac{1}{D_p(-p)} + \frac{1}{D_p(p)} \right) \frac{1}{D_\mu(p)} - \\
& -\frac{f^2(p^2)}{m_p^2} p^2 \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \left(\frac{1}{D_p(-p)} + \frac{1}{D_p(p)} \right) \frac{1}{D_\mu(p)} + \quad (12) \\
& + \left(\frac{f(p^2)}{m_p} \right)^2 2m_p p^2 \langle \gamma^\mu \gamma^\nu \rangle \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \left(\frac{1}{D_p(-p)} - \frac{1}{D_p(p)} \right) \frac{1}{D_\mu(p)} + \\
& + \left(\frac{f(p^2)}{m_p} \right)^2 2m_p p^0 \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \langle \gamma^\mu \hat{p} \gamma^\nu \rangle \left(\frac{1}{D_p(-p)} - \frac{1}{D_p(p)} \right) \frac{1}{D_\mu(p)},
\end{aligned}$$

where

$$D_p(\pm p) = p^2 \pm 2m_p p_0 + i\varepsilon, \quad D_\mu(p) = p^2 + 2m_\mu p_0 + i\varepsilon.$$

To make numerical calculations let transform (12) to convenient form. First of all we pick out in proton factor $1/D_p(-p)$ of crossed two-photon diagram the separate term with $\delta(p_0)$ function, which cancelled in future calculations by similar addendum of iteration part of quasipotential (3):

$$\frac{1}{D_p(-p)} = \frac{1}{2(m_p + m_\mu)} \left[-2\pi i \delta(p_0) - \frac{1}{p_0 + i\varepsilon} - \frac{D_\mu(p)}{(-p_0 + i\varepsilon) D_p(-p)} \right]. \quad (13)$$

It is useful also to represent the other proton factor $1/D_p(p)$, appearing in (12), in the following manner:

$$\frac{1}{D_p(p)} = \frac{1}{2(m_p - m_\mu)(p_0 + i\varepsilon)} \left[1 - \frac{D_\mu(p)}{D_p(p)} \right]. \quad (14)$$

Accounting these relations, we may transform different parts of (12) as follows:

$$\frac{p_0}{D_\mu(p)} \left[\frac{1}{D_p(-p)} - \frac{1}{D_p(p)} \right] = \frac{m_p}{m_p^2 - m_\mu^2} \left[\frac{1}{D_p(p)} - \frac{1}{D_\mu(p)} \right], \quad (15)$$

$$\frac{p_0}{D_\mu(p)} \left[\frac{1}{D_p(-p)} + \frac{1}{D_p(p)} \right] = \frac{m_\mu}{m_p^2 - m_\mu^2} \left[\frac{1}{D_\mu(p)} - \frac{1}{D_p(p)} \right], \quad (16)$$

$$\frac{p^2}{D_\mu(p)} \left[\frac{1}{D_p(-p)} + \frac{1}{D_p(p)} \right] \approx \frac{2}{m_p^2 - m_\mu^2} \left[\frac{m_p^2}{D_p(p)} - \frac{m_\mu^2}{D_\mu(p)} \right], \quad (17)$$

After that we rotate the p_0 contour to the imaginary axis and call the new variable ξ . The integrated factors are modified as follows:

$$\frac{1}{p^2 + i\varepsilon} \rightarrow -\frac{1}{\bar{p}^2 + \xi^2}, \quad \frac{1}{D_p(p)} \rightarrow \frac{-\bar{p}^2 + \xi^2 - 2im_p\xi}{(\bar{p}^2 + \xi^2)^2 + 4m_p^2\xi^2}. \quad (18)$$

As a result of such transformations we obtain that the contribution of 1, 3 and 4 form factor terms in (12) to muonic hydrogen HFS with the account of hadronic vacuum polarization is determined by triple integral of the following form:

$$\Delta E_1^{hf.s} = \tilde{E}_F \frac{m_p m_\mu}{m_p^2 - m_\mu^2} \frac{32\alpha^2}{\pi^2} \int_{s_0}^{\infty} \rho(s) ds \int_0^{\infty} d\xi \int_0^{\infty} p^2 dp \quad (19)$$

$$\frac{[\rho^2 - \frac{f^2}{m_p^2}(\bar{p}^2 + \xi^2)](3\xi^2 + 2\bar{p}^2) + 4\bar{p}^2 \rho f}{(\bar{p}^2 + \xi^2)(\bar{p}^2 + \xi^2 + s)} \left[\frac{m_p^2}{(\bar{p}^2 + \xi^2)^2 + 4m_p^2\xi^2} - \frac{m_\mu^2}{(\bar{p}^2 + \xi^2)^2 + 4m_\mu^2\xi^2} \right].$$

To do integration over s , we have used the parameterization of $\rho(s)$ by means of new experimental data on annihilation cross section $e^+e^- \rightarrow \text{hadrons}$ [10, 11, 12, 13]. The main contribution to the cross section σ^h is determined by the process $e^+ + e^- \rightarrow \pi^+ + \pi^-$. The cross section of this reaction is proportional to the squared module of π meson form factor F_π . Supposing that the ρ exchange diagram gives the basic contribution to F_π , we have used the following well-known expression of pionic form factor, taking into account the contribution of $\rho - \omega$ interference term [10, 13]:

$$F_\pi(s) = \frac{m_\rho^2(1 + d\Gamma_\rho/m_\rho)}{(m_\rho^2 - s + im_\rho\Gamma_\rho(p/p_0)^3 m_\rho/\sqrt{s})} + \zeta e^{i\phi} \frac{m_\omega^2}{m_\omega^2 - s + im_\omega\Gamma_\omega}, \quad (20)$$

where the ρ meson decay width $\Gamma_\rho=0.15$ GeV, ρ meson mass $m_\rho=0.77$ GeV and $d=0.5$. The effects of $\rho - \omega$ interference are described by the parameters from [10]. Substituting (20) to the spectral function

$$\rho_{\pi\pi}(s) = \frac{(s - 4m_\pi^2)^{3/2}}{12s^{5/2}} |F_\pi(s)|^2, \quad (21)$$

we obtain from (19) the corresponding contribution to HFS of muonic hydrogen:

$$\Delta E_{h1}^{hf.s}(\rho \rightarrow 2\pi) = 402.40 \text{ MHz}, \quad \tilde{E}^F = 15796.2 \text{ GHz}. \quad (22)$$

The term of (19), proportional to f^2/m_p^2 leads to the negative contribution 23,84 MHz in the energy spectrum. It is interesting to compare the quantity (19) with the result of its calculation in the point-like proton approximation, when we assume $\rho(k^2) \approx \rho(0) = e$, $f(k^2) \approx f(0) = e\kappa$. Keeping in mind that the main contribution in (19) is determined by first and third terms of (12), let transform their quantity in the hyperfine splitting of (μp) , fulfilling analytical integration over loop momentum p by means of Feynman parameterization. As a result we find:

$$\Delta E_h^{hf.s} = \left(\frac{\alpha}{\pi}\right)^2 \frac{m_\mu m_P}{m_P^2 - m_\mu^2} \tilde{E}_F \int_{4m_\pi^2}^\infty ds \rho(s) \int_0^1 dx [S(m_P, x) - S(m_\mu, x)], \quad (23)$$

where

$$S(m, x) = \frac{16 - 6x - x^2 + 4\kappa(8 - 6x + x^2)}{x^2 + \frac{s(1-x)}{m^2}}, \quad \tilde{E}^F = 8\alpha^4 m_\mu^2 m_P^2 / (m_P + m_\mu)^3.$$

Integration over parameter x may be done analytically also, and to integrate over s , we have used parameterization of $\rho(s)$ in the form (21). Then the contribution of first and third terms of (12) to HFS in the point-like proton approximation is equal to:

$$\Delta E_h^{hf.s}(\rho \rightarrow 2\pi) = 882.95 \text{ MHz}. \quad (24)$$

Comparing this value with the result (22), we may state that using of form factors (10), (11) in vertex operator (8) leads to essential expected decrease of corresponding correction by comparison to approximation $\rho(p^2) \approx \rho(0)$, $f(p^2) \approx f(0)$. The other parts of (12) give the contribution to HFS, which may be represented as a sum of two integrals similar to (19):

$$\Delta E_2^{hf.s} = \tilde{E}_F \frac{m_p m_\mu}{m_p^2 - m_\mu^2} \frac{32\alpha^2}{\pi^2} \int_{s_0}^\infty \rho(s) ds \int_0^\infty d\xi \int_0^\infty p^4 dp \quad (25)$$

$$\frac{2f^2(p^2)}{(\bar{p}^2 + \xi^2 + s)} \left[\frac{1}{(\bar{p}^2 + \xi^2)^2 + 4m_\mu^2 \xi^2} - \frac{1}{(\bar{p}^2 + \xi^2)^2 + 4m_p^2 \xi^2} \right],$$

$$\Delta E_3^{hf.s} = \tilde{E}_F \frac{m_p m_\mu}{m_p^2 - m_\mu^2} \frac{192\alpha^2}{\pi^2} \int_{s_0}^{\infty} \rho(s) ds \int_0^{\infty} d\xi \int_0^{\infty} p^2 dp \quad (26)$$

$$\frac{\rho(p^2) f(p^2) m_\mu (\bar{p}^2 + \xi^2)}{m_p (\bar{p}^2 + \xi^2 + s)} \left[\frac{1}{(\bar{p}^2 + \xi^2)^2 + 4m_\mu^2 \xi^2} - \frac{1}{(\bar{p}^2 + \xi^2)^2 + 4m_p^2 \xi^2} \right].$$

We have represented in the table the total contribution of all form factor terms of (12) with function $\rho_{\pi\pi}(s)$ to the HFS of muonic hydrogen, which is formed by the expressions (19), (25) and (26). Let take into account the contributions of ω and ϕ mesons to hyperfine structure of (μp) . To do this would require Breit-Wigner representation [13] of spectral function $\rho(s)$:

$$\rho_{\omega,\phi} = \frac{m_{\omega,\phi} \Gamma_{\omega,\phi}}{4s f_{\omega,\phi} [4(\sqrt{s} - m_{\omega,\phi})^2 + \Gamma_{\omega,\phi}^2]}, \quad f_{\omega,\phi} = \frac{\alpha^2 m_{\omega,\phi}}{12\Gamma_{ee}}, \quad (27)$$

where $m_\omega = 0.782$ Gev, $m_\phi = 1.019$ Gev, $\Gamma_\omega = 8.43$ Mev, $\Gamma_\phi = 4.43$ Mev, $\Gamma_\omega^{ee} = 0.60$ Kev, $\Gamma_\phi^{ee} = 1.37$ Kev [14]. Then after calculation of triple integrals (19) we obtain the summary contribution of the following kind:

$$\Delta E_h^{hf.s}(\omega) = 45.74 \text{ MHz}, \quad (28)$$

$$\Delta E_h^{hf.s}(\phi) = 48.97 \text{ MHz}. \quad (29)$$

Spectral density $R(s)$ was parameterized in the energy range $0.64 \leq s \leq 1.44$, between ω and ϕ resonances on the basis of present experimental data [12] on e^+e^- annihilation cross section in the form: $R(s) = A + Bs + Cs^2$ ($A=22.68$, $B=-37.72$, $C=15.84$). To obtain the contributions of J/Ψ and Υ particle families we have also used the Breit-Wigner formula (27) with known experimental values for total and partial decay widths of vector resonances to e^+e^- pair [14]. The calculation of background contribution at $s \geq 1.44$ was performed as in the paper [15]. The full integration region over s was divided on six intervals. In each interval the spectral density was founded by fitting of experimental points after subtraction of resonance contribution. Our numerical calculations show, that the main part of hadronic vacuum polarization contribution to hyperfine splitting of muonic hydrogen caused by π meson form factor (20). Theoretical error, not exceeding 10%, reflects spread in the choice of fitting parameters for cross section $\sigma(e^+e^- \rightarrow \text{hadrons})$ and for nucleon form factors. So, we actually see that whereas for hydrogen atom the relative contribution of hadronic vacuum polarization to ground state hyperfine splitting is about 10^{-8} [4], for muonium - $5.6 \cdot 10^{-8}$ [9, 15], then in the

case of muonic hydrogen it increases to $19.7 \cdot 10^{-6}$. The essential modification of hadronic vacuum polarization relative contribution to HFS of muonic hydrogen is connected with increase of particle relative motion momentum, which achieves the values of order of hadronic masses. Obtained in this paper value of correction $\Delta E_h^{hf.s} = 861.05 \text{ MHz}$, must be accounted for future comparison with experimental result and for more exact determination of δ^S and δ^P [5].

When constructing two-photon interaction quasipotential we take into account in the virtual Compton scattering amplitudes only elastic proton intermediate states. We have investigated exactly this mechanism of the hadronic vacuum polarization contribution to HFS of muonic hydrogen, which gives the correction to δ^S . It is possible additional "nonelastic" process considering the excitement in the intermediate states of "ladder" diagrams of Δ isobar or other nucleon resonances (correction δ^P). The contribution of such nonelastic mechanism demands special study.

Interval $\sqrt{s}$, Gev	$\Delta E_h^{hf.s} (\mu p)$, MHz
ρ	442.97
ω	45.74
ϕ	48.97
J/Ψ family	11.04
Υ family	0.11
Background	
$0.8 \leq \sqrt{s} \leq 1.2 \quad R = A + Bs + Cs^2$	131.90
$1.2 \leq \sqrt{s} \leq 3.0 \quad R = 2.843s^{-0.181}$	104.01
$3.0 \leq \sqrt{s} \leq 7.4 \quad R = 1.293s^{0.692}$	70.78
$7.4 \leq \sqrt{s} \leq 12.0 \quad R=3.91$	3.46
$12.0 \leq \sqrt{s} \leq 36.0 \quad R=4.01$	1.94
$36.0 \leq \sqrt{s} \leq 47.0 \quad R=4.13$	0.11
$\sqrt{s} \geq 47.0 \quad \text{QCD}$	0.02
Summary contribution	861.05

Nevertheless, it is possible to give the following approximate appreciation of hadronic vacuum polarization contribution to HFS of muonic hydrogen, taking into account proton polarizability. In the case of ordinary hydrogen the polarization correction δ^P in (2) is no more than 4 ppm [4]. It is equal approximately to 10% from elastic proton contribution δ^S . Then we can

obtain the approximate value of discussed contribution from proton polarizability replacing electron mass m_e by muon mass m_μ (as it follows from (23)) and adding one degree of α for calculation of vacuum polarization. So, in the case of muonic hydrogen such "nonelastic" mechanism may give contribution to HFS of order of $4 \text{ ppm} \frac{m_\mu}{m_e} \alpha \approx 6.0 \text{ ppm}$. This value is about 30% from relative contribution of hadronic vacuum polarization, obtained in this paper and equal to 19.7 ppm .

We are grateful to F. Jegerlehner, A. Karimkhodzaev, S. Karshenboim and V. Saleev for useful discussions. This work was supported by the Russian Foundation for Basic Research (grant no. 96-02-17309).

References

- [1] Bodwin G.T., Yennie D.R. //Phys. Rev. 1988. D37. N2. P.498.
- [2] Dvoeglazov V.V., Faustov R.N., Tukhtyaev Yu.N. //Fiz. Elem. Chast. i Atom. Yad. 1994. V.25. N1. P.144.
- [3] Bodwin G.T., Yennie D.T. //Phys. Rep. 1978. V.43. N6. P.267.
- [4] Karshenboim S.G. //Phys. Letters. 1997. A225. P.97-106.
- [5] Bakalov D., et al. //Proc. III Internat. Sympos. on Weak and Electromagnetic Interactions in Nuclei (WEIN-92) Dubna, Russia, 1992, P.656-662.
- [6] Faustov R.N. //Fiz. Elem. Chast. i Atom. Yad. 1972. V.3. N3. P.238.
- [7] Faustov R.N., Martynenko A.P. //Yad. Phys. 1987. V.45. P.770.
- [8] Berestetskii V.B., Lifshitz E.M., Pitaevskii L.P. Quantum Electrodynamics Nauka, Moscow, 1980.
- [9] Sapirstein J.R., et al. //Phys. Rev. 1984. D29. N10. P.2290.
- [10] Martinovic L., Dubnicka S. //Phys. Rev. 1992. D42. P.88
- [11] Swartz M.L. //Phys. Rev. 1996. D53. P.5268.
- [12] Eidelman S., Jegerlehner F. //Z.Phys. 1995. C67. P.585;
Jegerlehner F. //Preprint DESY-96-121, 1996.

- [13] Isaev P.S. Quantum Electrodynamics in High Energy Region, M., Energoatomizdat, 1984.
- [14] Review of Particle Properties //Phys. Rev. 1996. D54. N1. Part 1.
- [15] Faustov R.N., Karimkhodzaev A. //Yad. Phys. 1991. V.53. N4. P.1012.